

Leptons and Photons

L. Okun

ITEP, Moscow, 117259, Russia

Preprint ITEP No. 68-95

February 1, 2008

Abstract

It is assumed that three lepton families (ν_e, e) , (ν_μ, μ) , (ν_τ, τ) carry charges, which are sources of electronic, muonic and tauonic massless vector particles, respectively. Various manifestations of these hypothetical photons are discussed.

PACS: 14.80; 14.90

Keywords: electronic photons, muonic photons, tauonic photons.

We know six leptons: three electrically charged ones, e, μ, τ , and three electrically neutral ones, ν_e, ν_μ, ν_τ . We know only one photon that is coupled to the electric charge of particles. One of the most fundamental properties of the electric charge is its conservation. There exist three different leptonic quantum numbers: electronic for e, ν_e , muonic for μ, ν_μ , and tauonic for τ, ν_τ , which at the present level of our ignorance are conserved. If they are strictly conserved they may serve as corresponding charges emitting and absorbing $\gamma_e, \gamma_\mu, \gamma_\tau$ – electronic, muonic and tauonic photons, respectively. These hypothetical charges and hypothetical photons and their possible manifestations are the subject of this letter.

If neutrino oscillations are discovered experimentally, that will prove that the number of leptonic charges is reduced. Observation of transitions $\nu_e \leftrightarrow \nu_\mu \leftrightarrow \nu_\tau$ would leave open only one window – that for the existence of total leptonic charge. In a similar manner observation of neutrinoless double beta decay ($2n \rightarrow 2e + 2p$) would mean that electronic charge is definitely

not conserved and hence there is no electronic photons. (A non-conserved charge cannot be a source of massless vector particles and of a Coulomb-like potential, behaving as $1/r$). The non-conservation of electronic or muonic charge would be also proved by the relevant proton decays, promised to us by the grand unified theories. (Note that in some of them B-L – the difference between baryonic and leptonic charges is conserved.) But today there is still plenty of room for speculations about all three types of leptonic photons.

It should be stressed that the existence of leptonic photons does not seem to solve any of the existing problems. In that sense they are not needed. But who needed muon when it was discovered?

The leptonic photons are not only not needed; they are undesirable, as the additional Abelian gauge symmetries connected with them would produce triangle anomalies, which without special care would destroy the renormalizability of the Standard Model.

After presenting all necessary warnings let us proceed. Analogously to the electric charge e and to the fine structure $\alpha = e^2/4\pi$, let us introduce g_e, g_μ, g_τ and $\alpha_e = g_e^2/4\pi$, $\alpha_\mu = g_\mu^2/4\pi$, $\alpha_\tau = g_\tau^2/4\pi$. One of our goals would be to establish the upper limits on $\alpha_e, \alpha_\mu, \alpha_\tau$.

For α_e a straightforward limit is given by the tests of independence of gravitational acceleration on the material of the probe body. These tests were started many years ago by Eötvös [1] at the accuracy 10^{-8} . More recent tests [2, 3] have reached 10^{-12} . The electronic "Coulomb force" should give repulsion, proportional to the product of the numbers of electrons in the earth (or sun) and the test body. Therefore the variation of the acceleration with the kind of the test body atoms would be [4, 5]

$$1 + \frac{\alpha_e}{\alpha_g} \left(\frac{\bar{Z}}{\bar{A}} \right) \left(\frac{Z}{A} \right)$$

where $\alpha_g = Gm_p^2 = 6 \cdot 10^{-39}$ (G – the Newton gravitational constant, m_p – mass of proton), $\bar{Z}$ and $\bar{A}$ are the numbers of electrons and nucleons in an average atom of the earth ($\bar{Z}/\bar{A} = 0.5$) or sun ($\bar{Z}/\bar{A} = 1$); while Z and A refer to an atom of a test body. If we compare two balls, one of copper, another of lead, the difference in acceleration would be

$$\frac{\alpha_e}{\alpha_g} \left(\frac{\bar{Z}}{\bar{A}} \right) \left(\frac{Z_{Cu}}{A_{Cu}} - \frac{Z_{Pb}}{A_{Pb}} \right) < 10^{-12}$$

Taking into account that $\frac{Z_{Cu}}{A_{Cu}} = 0.46$, $\frac{Z_{Pb}}{A_{Pb}} = 0.40$ we get

$$\alpha_e \leq 10^{-49} .$$

This upper limit is obtained under assumption that the electronic charge of the sun (or earth) is not neutralized by antileptons. Such neutralization is impossible if the only light (and electrically neutral) antileptons are antineutrinos [6, 7]. The number of $\bar{\nu}$'s cannot be equal to that of e 's inside a celestial body. Because of Pauli principle the momenta of $\bar{\nu}$'s are of the order of electron momenta. Thus their kinetic energy is much larger than that of electrons, while the potential energy is smaller by a factor $\alpha_e/\alpha \ll 1$. Thus, there can be no equilibrium between equal amounts of electrons and antineutrinos.

In principle, one can think [6], that the electronic charge of the earth is screened by some neutral leptonic bosons, carrying the electronic (leptonic) charge, such particles were discussed in ref. [8]. In ref. [9, 10, 7] they were assumed to be electronic antisneutrinos – $\tilde{\bar{\nu}}_e$ – supersymmetric scalar partners of $\bar{\nu}_e$ with $m_{\tilde{\bar{\nu}}_e} \leq 1$ eV. But for the Eötvös type upper limit to become irrelevant, the screening must be perfect. For instance, to allow $\alpha_e \sim 10^{-20}$ the screening must have accuracy 10^{-29} which does not seem plausible. Especially difficult to get a substantial neutralization of electronic charge in bodies with dimensions smaller than, say, 1 cm. Having some of their $\tilde{\bar{\nu}}'$ outside, these bodies may explode [6, 7]. The explosion would be caused by the leptonic repulsion of their unscreened electrons. Let us discuss the mechanism of explosion in some detail for the case of no screening. Consider a sphere of radius r with density n of atoms per cm^3 and number of electrons per atom Z . Omitting numerical coefficients we get for the force of the volume repulsion

$$F_R \sim \alpha_e \frac{(Znr^3)^2}{r^2} .$$

The force of attraction of the two halves of the sphere is proportional to the separation surface and can be estimated as

$$F_A = nr^2 a \frac{\alpha^2}{a^2} ,$$

where $a \sim 10^{-8}$ cm is the dimension of an atom, while α^2/a^2 is a crude estimate of the Van-der-Waals force. The sphere would become unstable at

$r > r_c$. The critical radius r_c is determined by the equation

$$F_R = F_A \ ,$$

wherefrom we obtain

$$r_c = \left(\frac{\alpha^2}{Z^2 n \alpha_e a} \right)^{1/2} \sim a \frac{\alpha}{Z \alpha_e^{1/2}} \ .$$

For $\alpha_e \sim 10^{-20}$ and $Z \sim 10$ we get $r_c \sim 1\text{mm}$. Further theoretical work is needed to prove or disprove the possibility of screening of the electronic charge of celestial bodies.

Let us now consider α_μ and α_τ . The upper limits for them can be derived from experiments with μ 's, τ 's, and from the neutrino experiments.

An upper limit on α_μ from the muon magnetic moment is determined by the experimental accuracy of

$$a_\mu \equiv \frac{1}{2}(g_\mu - 2) \ .$$

According to [11]

$$\frac{\Delta a_\mu^{exp}}{a_\mu} \simeq 10^{-5} \ .$$

While the virtual γ gives correction to a_μ close to $\alpha/2\pi$, the virtual γ_μ gives $\alpha_\mu/2\pi$. Hence α_μ must be at least five orders of magnitude smaller than α . This means that about 10^{-5} of all photons emitted by muons could be γ_μ 's. Muonic photons would be emitted more effectively by ν_μ 's, for instance in pion decays the ratio

$$\frac{\Gamma(\pi \rightarrow \mu \nu_\mu \gamma_\mu)}{\Gamma(\pi \rightarrow \mu \nu_\mu)} \sim \frac{2\alpha_\mu}{\pi} \frac{d\omega}{\omega} \ln \frac{2\Delta}{m_{\nu_\mu}} \ ,$$

where ω is the energy of the photon in the pion rest frame, $\Delta = m_\pi - m_\mu$, and we assume $\omega \ll \Delta \ll m_\mu$. Thus ν_μ -beams would be accompanied by γ_μ -beams. Muonic photons would be penetrating and would produce pairs of muons when colliding with nuclei. The cross-section of this process on a pointlike nucleus with charge Z at asymptotically high energy ω would be [12]:

$$\sigma = \frac{\alpha^2 Z^2 \alpha_\mu}{m_\mu^2} \cdot \frac{28}{9} \left[\ln \frac{2\omega}{m_\mu} - \frac{109}{42} \right] \ .$$

The search for such pairs may further decrease the upper limit on α_μ .

Similar limit on α_τ from $(g_\tau - 2)$ is non-existent. Therefore a search for τ -photons in τ - and ν_τ -experiments is even more interesting. If CHORUS and NOMAD do not find ν_τ -oscillations, they may find narrow $\mu^+\mu^-$ pairs from γ_μ 's or $\tau^+\tau^-$ pairs from γ_τ 's.

Cosmological limit on the mass density of relic ν_μ 's and ν_τ 's gives lower limits on α_μ and α_τ [4], assuming that $m_{\nu_\mu}, m_{\nu_\tau} \gtrsim 0.1 \text{ KeV}$ and that these neutrinos were not burned out by some exotic processes:

$$\alpha_\mu > 10^{-12} \left(\frac{m_{\nu_\mu}}{1 \text{ KeV}} \right) , \quad \alpha_\tau > 10^{-12} \left(\frac{m_{\nu_\tau}}{1 \text{ KeV}} \right) .$$

These lower limits follow from the requirement that the annihilation of the pairs $\nu_\mu \bar{\nu}_\mu$ and $\nu_\tau \bar{\nu}_\tau$ into leptonic photons was fast enough to guarantee an acceptable mass density of these particles.

Of special interest are the neutrino electric charges induced by the mixing of ordinary and leptonic photons [4]. The mixing appears through the vacuum polarization loops of e, μ, τ because these leptons carry both leptonic and ordinary charges. The simplest Lagrangian of four photons without mixing may be written (omitting the four-vector indices) in the form:

$$\mathcal{L} = \sum_i \left(\frac{1}{4} F_i^2 + g_i A_i j_i \right) ,$$

where $i = 0$ refers to ordinary photon γ (so that $g_0 \equiv e$), while $i = 1, 2, 3$ refer to e, μ, τ , correspondingly.

In the one-loop approximation three terms have to be added to $\mathcal{L}$: FF_e , FF_μ , FF_τ with coefficients proportional to $\sqrt{\alpha\alpha_e}$, $\sqrt{\alpha\alpha_\mu}$ and $\sqrt{\alpha\alpha_\tau}$, correspondingly. In the two-loop approximation three more terms should be added $F_e F_\mu$, $F_e F_\tau$, $F_\mu F_\tau$ with coefficients $\alpha\sqrt{\alpha_e\alpha_\mu}$, $\alpha\sqrt{\alpha_e\alpha_\tau}$ and $\alpha\sqrt{\alpha_\mu\alpha_\tau}$, correspondingly. Moreover the coefficients of the diagonal terms $\frac{1}{4}F^2$, $\frac{1}{4}F_e^2$, $\frac{1}{4}F_\mu^2$, $\frac{1}{4}F_\tau^2$ should be changed.

Were α_e not negligibly small, the induced electric charges, of ν_e and of e could be observable:

$$Q_{\nu_e} - Q_n = Q_e - Q_p \neq 0 .$$

As a result the electrical neutrality of atoms would be lost . (It is tested at the level of $10^{-20} - 10^{-23}$; for a collection of experimental data see e.g. ref. [13]).

The mixings $\gamma \leftrightarrow \gamma_\mu$ and $\gamma \leftrightarrow \gamma_\tau$ will not damage the neutrality of atoms, but will produce nonvanishing electric charges of ν_μ and ν_τ .

For light ν_μ and ν_τ ($m \ll 1$ KeV) the upper limits on their electric charges are given by the neutrino luminosity of the sun [14]

$$Q_{\nu_\mu} \quad \text{and} \quad Q_{\nu_\tau} \leq 10^{-13} \quad .$$

Independently of the value of the mass of ν_μ , an upper limit on its electric charge can be derived from the data on $\nu_\mu e$ -scattering [16]:

$$Q_{\nu_\mu} \leq 10^{-8} \quad .$$

The mixing coefficients, derived in one- or two-loop approximation are logarithmically divergent. This divergences are cancelled by corresponding counterterms. Thus the resulting mixing coefficients and the neutrino electric charges might be quite different from their naive estimates. After diagonalization of ten $F_i F_k$ terms we are left with four diagonal "photons", but each of them will be coupled to all four currents.

Further complications arise if we take into account the box-type muon or tauon loops [4] which contribute to the magnetic moments of charged particles. A lot of exotic phenomena (similar to neutrino oscillations) appear if leptonic photons are not strictly massless, but have tiny masses [16].

Finally, let us return to the problem of anomalies.

Looking for an anomaly-free scheme with leptonic photons, for which Eötvös type experiments are irrelevant, it is natural to consider a scheme with one leptonic photon which has no coupling to (e, ν_e) and has non-vanishing couplings of equal strength and opposite sign to (μ, ν_μ) and (τ, ν_τ) . In this case the triangle with one leptonic photon and two electroweak gauge fields vanishes due to cancellation between (μ, ν_μ) and (τ, ν_τ) , while the triangle with two leptonic photons and one electroweak gauge field vanishes due to cancellation between neutrino and its charged partner. In the framework of this scheme the leptonic photons emitted by muons in the high-energy neutrino experiments could in principle produce not only pairs $\mu^+ \mu^-$, but also pairs $\tau^+ \tau^-$. However the cross-section of the latter process would be three orders of magnitude smaller.

Another type of anomaly-free scheme has been suggested to me by M.B.Voloshin. In his scheme the baryo-leptonic photon is coupled to the difference of

baryonic and leptonic charges: B-L. Its coupling to all three generations has to be universal because of the Kobayashi-Maskawa mixing between quarks.

As was pointed out to me by A.D.Dolgov, recent observational data on light element abundances allow existence of one extra massless particle. (See e.g. B.Fields, K.Kainulainen, K.Olive, hep-ph 9512321.)

I am grateful to A.D.Dolgov for useful discussions of the manuscript of this letter. This research was made possible in part by grant 93-02-14431 of the Russian Foundation for Basic Research. The main content of this letter was presented at the Symposium in honour of Klaus Winter on the occasion of his 65th birthday, on December 6, 1995 at CERN.

References

- [1] R.V.Eötvös, D.Pekar, and E.Fekete, Ann. der Phys. **68** (1922) 11.
- [2] R.H.Dicke, G.Krotkov, and P.G.Roll, Ann. of Phys. **26** (1964) 442.
- [3] V.B.Braginsky and V.I.Panov, Sov. Phys. JETP **34** (1972) 463.
- [4] L.B.Okun, Yadernaya Fizika **10** (1969) 358 (in Russian).
Sov. Journal of Nucl. Phys. **10** (1969) 206 (in English).
- [5] L.B.Okun. Leptons and Quarks. North Holland. 1982, p. 166.
- [6] L.B.Okun. A remark on leptonic photons, June 1972 (Manuscript in Russian, presented at the ITEP theoretical seminar).
- [7] S.I.Blinnikov, A.D.Dolgov, L.B.Okun, and M.B.Voloshin. How Strong can the coupling of leptonic photons be? Preprint ITEP-31-95, TPI-MINN 95/12-T, UMN-TH-1342-95, hep-ph 9505444, to be published in Nucl. Phys. **B458** (1996) 5264.
- [8] L.B.Okun, ZhETF **47** (1964) 1777; Sov. Phys. JETP **12** (1964) 250.
- [9] A.K.Çiftçi, S.Sultansoi, and S.Türköz. Lepton charge as possible source of new force. Ankara University preprint AU/94-03 (HEP).
- [10] A.K.Çiftçi, S.Sultansoi, and S.Türköz. Phys. Lett. **B 355** (1995) 494.

- [11] Review of Particle Properties. Phys. Rev. **D50**, No. 3, part I (1994) 1396.
- [12] A.I.Akhiezer and V.B.Berestetskiy. Quantum Electrodynamics. Moscow, Nauka, 1969. §30.1.
- [13] L.B.Okun, M.B.Voloshin, and V.I.Zakharov, Phys. Lett. **138B** (1983) 115.
- [14] J.Bernstein, M.Ruderman, and G.Feinberg, Phys. Rev. **132** (1968) 1227.
- [15] CHARM II Collab. P.Vilain et al., Phys. Lett. **B281** (1992) 159.
- [16] L.B.Okun, ZhETF **83** (1982) 892 (in Russian);
Sov. Phys. JETP **56** (1982) 502 (in English).